Document downloaded from:

[\[http://redivia.gva.es/handle/20.500.11939/6243\]](http://redivia.gva.es/handle/20.500.11939/6243)

This paper must be cited as:

[Tena, A.; Nieves, E.; Herrero, J.; Urbaneja, A. (2018). Defensive behaviors of the new mealybug citrus pest, delottococcus aberiae (hemiptera: Pseudococcidae), against three generalist parasitoids. Journal of Economic Entomology, 111(1), 89-95.]

ivia
Institut Valencià
d'Investigacions Agràries

The final publication is available at

[\[http://dx.doi.org/10.1093/jee/tox307\]](http://dx.doi.org/10.1093/jee/tox307)

Copyright [Oxford University Press]

Article type: *Original Article*

Journal of Economic Entomology

Running title: Defensive behaviors of *Delottococcus aberiae*

Defensive behaviors of the new mealybug citrus pest, *Delottococcus aberiae* De Lotto, against three generalist parasitoids

Alejandro Tena¹, Eliana Nieves², Julieta Herrero¹, Alberto Urbaneja¹

¹Instituto Valenciano de Investigaciones Agrarias (IVIA), Unidad Asociada de Entomología Agrícola UJI-IVIA: Centro de Protección Vegetal y Biotecnología: CV-315, Km. 10,7 – 46113 Moncada (Valencia). Spain.

²Centro de Estudios Parasitológicos y de Vectores (CEPAVE), La Plata, Argentina

* Corresponding author: Dr. Alejandro Tena

Centro Protección Vegetal y Biotecnología.

Instituto Valenciano de Investigaciones Agrarias (IVIA)

Carretera Moncada-Náquera km 4.5.

46113 Moncada (Valencia, Spain)

Phone: (+34) 96 342 41 51

Fax: (+34) 96 342 40 01

Email: atena@ivia.es

Abstract: *Delottococcus aberiae* De Lotto (Hemiptera: Pseudococcidae) is an invasive mealybug that has become a citrus pest in Europe. This mealybug species causes serious damage because it deforms the fruits. Here, we studied the defensive behavior of *D. aberiae* when it was attacked by three parasitoid species: *Acerophagus angustifrons* (Gahan), *Anagyrus* sp. near *pseudococi* (Girault) and *Leptomastix algerica* Trjapitzin (Hymenoptera: Encyrtidae). *Anagyrus* sp. near *pseudococi* and *L. algerica* detected and accepted nymphs and adult females of *D. aberiae* whereas *A. angustifrons* only accepted adults. We recorded four defensive responses of *D. aberiae* to parasitoid attacks: abdominal flipping, swiveling around the inserted stylet, withdrawing the stylet and walking away and, occasionally, they secreted ostiolar fluids. Despite these defensive behaviors, the mealybug did not escape parasitism from any of the tested parasitoids, even though *A. angustifrons* needed more than 15 min to parasitize. We also analyzed the nutritional value of the honeydew excreted by *D. aberiae* for *A. angustifrons* and *A. sp. near pseudococi*. Females and males of these parasitoids lived more than 28 days when fed sucrose, but they lived fewer than three days when fed *D. aberiae* honeydew. Therefore, *D. aberiae* excretes honeydew of poor quality for parasitoids. The consequences of these biological traits of *D. aberiae* for its biological control are discussed.

Keywords: *Acerophagus*, *Anagyrus*, *Leptomastix*, biological control, honeydew, nutritional ecology

INTRODUCTION

Delottococcus aberiae (De Lotto) (Hemiptera: Pseudococcidae) is a polyphagous mealybug native to sub-Saharan Africa (Miller and Giliomee 2011; Beltrà et al. 2015b), which was detected for the first time in 2009, as a citrus pest in the Mediterranean basin (Beltrà et al. 2013). After a comprehensive but unsuccessful eradication program, this mealybug established as a serious pest in the region of Les Valls (Valencia, Eastern Spain), which is located in the core of the main citrus producing area of the Mediterranean basin (Beltrà et al. 2013, Tena et al. 2014). *Delottococcus aberiae* increases its populations during spring and nymphs and adults settle and feed on the surface of young fruitlets under the calyces (Martínez-Blay et al. 2017). This feeding habit causes reduction and distortion of growing fruit (Pérez-Rodríguez 2017). Encyrtid parasitoids and predatory coccinelids are among the most successful families of natural enemies used in biological control of mealybugs (Barlett 1978, Moore 1988). Numerous biological control programs have been carried out and, as result of these efforts, most pseudococcids are under excellent biological control (Moore 1988, Franco et al. 2004). Some genera of the Encyrtidae such as *Acerophagus* or *Leptomastix* have been used successfully in the biological control of other mealybug species, including *Phenacoccus peruvianus* Granara de Willink, *Phenacoccus herreni* Cox & Williams, and *Paracoccus marginatus* Williams and Granara de Willink (van Driesche et al 1987, Muniappan et al. 2006, Beltrà et al. 2015). Parasitoids of the Anagyrini tribe, which contains the genus *Anagyrus*, have been cited as primary parasitoids of *Planococcus* mealybugs as *P. citri* (Risso) and *P. ficus* Signoret (Noyes & Hayat 1994, Franco et al. 2004). The success of mealybug parasitoids depends, in part, on their capacity to overcome their hosts' defenses (Bartlett 1978, Gross 1993, Blumberg 1997, Blumberg and van Driesche 2001, Daane et al. 2007, Campos-Rivela 2008, Tena et al. 2012, Sime and Daane 2014, Beltrà et al. 2015a, Bugila et al. 2015). In the case of *D. aberiae*, we have recently demonstrated that it is able to encapsulate the eggs of three generalist parasitoids of mealybugs: *Acerophagus angustifrons* (Gahan), *Anagyrus* sp. near *pseudococi* (Girault) and *Leptomastix algerica* Trjapitzin (Hymenoptera: Encyrtidae) (Tena et al. 2017). However, the

behavioral defenses of this new citrus pest against parasitoids have not been described. These parasitoids were selected because they are native or naturalized species in Spain, are considered generalist parasitoids of pseudococcids (Universal Chalcidoidea Database 2016, Martínez-Ferrer 2003, Campos-Rivela 2008), and are able to parasitize *D. aberiae* (Tena et al. 2017).

Despite being vulnerable to parasitoids, mealybugs have developed diverse defensive strategies that can hinder biological control (Blumberg 1997, Gullan and Kosztarab 1997, Gutierrez et al. 2008). They hide from their natural enemies by adopting cryptic behavior (Vet and Dicke 1992, Gross 1993, Foldi 1997, Gutierrez et al. 2008). When these defense mechanisms fail, mealybugs carry out evasive movements similar to those performed by aphids (Hemiptera: Aphididae) and soft scales (Hemiptera: Coccidae) (De Farias and Hopper 1999, Wyckhuys et al. 2008, Tena et al. 2012). Some of these movements, such as wriggling, swiveling around the stylet, getting up, walking away, and secreting defensive exudates, have been recorded to thwart parasitoid oviposition (Bynum 1937, Boavida et al. 1995, Bokonon-Ganta et al. 1995, Cadée and van Alphen 1997, Bugila et al. 2014, Beltrà et al. 2015a). The elucidation of such interactions is crucial to understanding the potential of these parasitoids and to select parasitoids for biological control.

Mealybugs excrete large amount of honeydew that can be used by their parasitoid as a carbohydrate source. Generally, honeydew has been considered a poorer carbohydrate source for adult parasitoids when compared to nectar (Wäckers et al. 2008, Tena et al. 2016). However, honeydew is available in citrus and the honeydew of at least two mealybug species significantly increases the longevity and fecundity of different parasitoids (Avidov et al. 1970, Tena et al. 2013). In the case of *D. aberiae*, it is unknown whether its honeydew is suitable for parasitoids. In order to design future biological control programs it is important to determine the sugar sources available and its quality for parasitoids (Tena et al. 2016).

The aims of this study were: i) to compare the host selection behavior of the parasitoids *A. angustifrons*, *A. sp. near pseudococi* and *L. algerica* in relation to the mealybug *D. aberiae*; ii) to describe the defensive behavior of *D. aberiae* against these parasitoids, and iii) to analyse the

suitability of *D. aberiae* honeydew as a carbohydrate source for these parasitoids. These results will increase our knowledge of the defensive mechanisms of this new citrus pest and whether additional carbohydrates are necessary to efficiently rear parasitoids in the insectary or conserve their populations in the field.

MATERIAL AND METHODS

Mealybug rearing and parasitoid sources

The colony of *D. aberiae* was established in the laboratory of Entomology at Instituto Valenciano de Investigaciones Agrarias (IVIA) using specimens collected in a citrus orchard located in Quartell (Les Valls, Valencia, Spain) in 2013. Mealybugs were reared on organically ripen lemons inside cardboard boxes (30 × 22 × 25 cm) in which egg cartons were placed in the bottom. Approximately 65% of the surface of each lemon was covered with red paraffin around the mid-section to retard its desiccation. The red paraffin was prepared with a mixture of 1 kg of paraffin pearls (Parafina USP Perlas, Guinama S.L., Alboraya, Spain) and 1 g of red pigment (Sudan III, Panreac Química S.A., Castellar del Vallés, Spain) following the same procedure used for the rearing of California red scale, *Aonidiella aurantii* (Maskell) (Hemiptera: Diaspididae) (Vanaclocha et al. 2012). Every two weeks, four or five fresh lemons were introduced within each box and the dried ones were removed. The mealybug colony was maintained in darkness in a climatic chamber at 25 ± 1 °C and 70 ± 5% RH.

The parasitoids *A. angustifrons*, *A. sp.* near *pseudoccoci* and *L. algerica* were obtained from Koppert Biological Systems S.L. (Águilas, Murcia, Spain). For the behavioral assay, the bottles with the pupae were opened and introduced in wooden and crystal rearing boxes (51 × 51 × 41 cm) with holes covered with mesh in the wall. Honey drops on the walls and an Eppendorf tube filled with water and a piece of cotton were provided. Rearing boxes were kept in the laboratory at room temperature until parasitoids were used. Emerged parasitoids were collected daily at 15:00, sexed and individualized in a 3.0 × 0.8 cm in diameter glass and sealed with cotton. Parasitoids could mate during the 24 hours that both sexes were together in the rearing boxes. A

drop of honey was provided on the wall. Female parasitoids were used 2-4 days later in the behavioral assay.

For the survival assay, parasitoid pupae were introduced in a 3.0 x 0.8 cm diameter glass vials sealed with a piece of muslin mesh. At emergence, parasitoids were sexed and diets were provided and replaced every two days as explained below. Parasitoids were kept in a climatic chamber at 25 ± 1 °C, $70 \pm 5\%$ RH and a photoperiod of 14L: 10D h in both assays.

Parasitoid oviposition behavior and host defense responses

The oviposition behavior of *A. angustifrons*, *A. sp.* near *pseudoccoci* and *L. algerica* were evaluated by direct observations, following methods similar to those described by Desneux et al. (2009). We also observed *D. aberiae* defensive behaviors and the effectiveness of these behaviors in preventing parasitism. Finally, we assessed the relationship between *D. aberiae* defensive behaviors and oviposition success of the three parasitoids.

The substrate for observations was the surface of a lemon under a binocular microscope illuminated with cold light. We placed one individual of *D. aberiae* on lemon surface using a fine brush and allowed it to establish for 24 hours. The individual of *D. aberiae* was either a 3rd instar or a preovipositing adult female, which are the preferred instars for these parasitoids (Tena et al. 2017). Then, a glass vial with a parasitoid female was opened and placed near the host mealybug on the lemon surface. Observations began when the vial came into contact with the lemon and ended when: i) oviposition finished (parasitism), ii) the female contacted and rejected the host (rejection) or iii) the female did not contact the host after 15 minutes (indifference). The frequency and time spent by the parasitoid and the host in each behavioral event were recorded by voice recording with the program Audacity®, the Free, Cross-Platform Sound Editor (<http://www.audacityteam.org/>). The time spent to find the host was also recorded. We recorded parasitoid behaviors following Beltrà et al. (2015a): i) antenna drumming—drumming the host body with the antennae, ii) ovipositor tap —assessing the host by tapping the

body with the ovipositor, iii) oviposition drill that included probing the host's body with the ovipositor and abdominal movements with the ovipositor inside the host's body, and iv) remove ovipositor—removing ovipositor.

Mealybug defensive action patterns were classified following Boavida et al. (1995) and Bugila et al. (2014): i) abdominal flipping—refers to repeated up and down movements of the hind half of the mealybug, ii) moving—swiveling around the inserted stylet, iii) reflex bleeding— refers to the secretion of ostiolar fluid, iv) walking away—the mealybug may also escape from the aggressor. We considered that a mealybug defended itself from an oviposition attempt when it carried out at least one of the above behaviors.

We also recorded the host detection and acceptance rates. We considered that a host was detected and accepted when the parasitoid drummed the mealybug with its antenna, and inserted its ovipositor in an attempt to deposit an egg, respectively. To confirm the oviposition, the parasitoid was removed and the mealybug was carefully transferred to a tender leaf of citrus clementine (*Citrus sinensis* cv. clemenules) after the observation period. The leaf was introduced over a layer of bacteriological agar (20 g/l) with the petiole inserted in the agar within a 5.3-cm diameter Petri dish with a 3-cm diameter hole covered by a muslin mesh to permit ventilation (Tena et al. 2017). The arenas were placed undisturbed in a climatic chamber (25 ± 1 °C, $70 \pm 5\%$ RH and a photoperiod of 14L: 10D h). Ten days later we confirmed whether or not the mealybugs were parasitized (i.e. mummified or contained encapsulated parasitoid eggs) (Tena et al. 2017). Observations were replicated between 12 and 24 times depending on the parasitoid species and host instar.

Effect of *Delottococcus aberiae* honeydew on parasitoid longevity

To assess the quality of the honeydew excreted by *D. aberiae* as a sugar source for *A. sp.* near *pseudococci* and *A. angustifrons*, we compared their longevity when they fed on 2 M (mol/L) sucrose (catalog number S0389, Sigma–Aldrich®, UK), honeydew and water. Female and male parasitoids were tested in this assay. Water was supplied for all treatments by spraying the vials

through the muslin mesh. Honeydew and sucrose (three droplets of 5 μ l) were provided *ad libitum* on 0.5 cm² piece of Parafilm[®] (Tena et al. 2013). Honeydew was collected over a period of 24 h by placing pieces of Parafilm[®] below a colony of mealybugs settled on lemons fruits in the climatic chambers. The presence of honeydew on the pieces of Parafilm[®] was checked under a binocular and then kept at -20 °C within a Petri dish until further use (Hogervorst et al. 2007, Tena et al. 2013). Vials with parasitoids were checked daily from 8:00 to 10:00 to determine the number of surviving parasitoids. Accidental deaths (wasps stuck in honey) were not used in the data analyses. Between 15 and 40 parasitoids were used per treatment.

Statistical analysis

We applied generalized linear modeling (GLM) techniques assuming Poisson and binomial error variance to construct models using the below-mentioned dependent variables and parasitoid species as the explanatory variable. Initially, we assumed a Poisson error variance for the count variable of “defensive responses per host” and a binomial error variance for the proportional variable of host detection and acceptance. When an over- or underdispersion was detected, we re-evaluated the significance of the explanatory variables using an F-test after rescaling the statistical model by a Pearson’s chi-square divided by the residual degrees of freedom (Crawley 2007). The data are presented as the means of untransformed proportion and count data (in preference to less intuitive statistics such as the back-transformed means of logit-transformed data). For the variable “oviposition time”, we assumed normal distribution and an ANOVA was carried out with parasitoid as explanatory variable. Moreover, Tukey’s post hoc tests were performed. The effect of the feeding treatments on the longevity of adult parasitoids was represented by Kaplan–Meier survivorship curves and analyzed by log-rank tests. All the analyses were performed by means of statistical software R (<http://www.R-project.org>) and the packages “OIsurv” for the survival analysis and “MULTCOMP” for the Tukey’s post hoc tests.

RESULTS

Parasitoid oviposition behavior and host defense responses

Host detection and acceptance

Detection rates of 3rd instar nymphs of *D. aberiae* by the three parasitoids ranged between 0.23 and 0.94 (Table 1). The rate of host detection in *A. sp. near pseudococci* was significantly higher than in *A. angustifrons* and *L. algerica* ($\chi^2 = 49.3$, $df = 2$, 50 , $P < 0.0001$). In the case of *D. aberiae* adult females, detection rates ranged between 0.67 and 0.88, but no significant differences were found among the three parasitoid species ($\chi^2 = 58.14$, $df = 2$, 52 , $P = 0.22$). *Anagyrus sp. near pseudococci* and *L. algerica* accepted a significantly higher proportion of 3rd instar nymphs than *A. angustifrons* ($\chi^2 = 53.65$, $df = 2$, 50 , $P = 0.0002$) (Table 1). In this last parasitoid species only one female of *A. angustifrons* parasitized a 3rd instar nymph. In the case of *D. aberiae* adult females, no significant differences were found among the three parasitoid species ($\chi^2 = 62.7$, $df = 2$, 52 , $P = 0.07$).

Oviposition behavior

Anagyrus sp. near pseudococci and *L. algerica* needed about ~90 sec to parasitize 3rd instar nymphs of *D. aberiae* and there were not significant differences between the two species ($F_{1, 18} = 0.40$, $P = 0.53$). These parasitoids spent ~65 sec to parasitized adults of *D. aberiae* and, again, there were not significant differences between the two species ($F_{1, 23} = 2.45$, $P = 0.13$). They spent most of this time drilling and laying the eggs (Fig. 1). In the few cases in which *A. angustifrons* accepted adult mealybugs of *D. aberiae* as hosts, females needed more than 15 min (1048-1489 sec) to parasitize adult mealybugs. Similarly, they spent most of this time drilling and laying the eggs (84.26%-98.46% of the total time spent).

Host defence

Overall, 92% (23 out of 25) and 88.6% (31 out of 35) of the nymphs and adults of *D. aberiae* defended themselves from parasitoid attacks, respectively. Third instar *D. aberiae* nymphs showed a mean of 5.79 ± 1.1 defensive responses against *A. sp. near pseudococci* and 3.0 ± 0.69 against *L. algerica* ($F_{1, 23} = 4.04$, $P = 0.056$). *Delottococcus aberiae* adults showed a mean of 5.8 ± 2.27 defensive responses against *A. angustifrons*, 6.29 ± 1.12 against *A. sp. near pseudococci*

and 4.56 ± 1.48 against *L. algerica* ($F_{2,32} = 0.43$, $P = 0.65$) (Fig. 2). The most common defensive response of *D. aberiae* during the oviposition process was abdominal flipping (Fig. 2). Mealybugs also defended themselves by swiveling around the inserted stylet and by withdrawing the stylet and walking away (Fig 2). Finally, some individuals also secreted ostiolar fluids during the parasitoid oviposition. However, these defensive responses did not succeed and only three mealybugs out of the 54 escaped parasitism (one defended against *A. angustifrons* and two against *A. sp. near pseudococci*).

Effect of *Delottococcus aberiae* honeydew on parasitoid longevity

The diet provided to the parasitoids significantly influenced the life span of both parasitoids (*A. angustifrons* females: (Log-rank test) $\chi^2 = 43.45$, $df = 2$, $P < 0.001$, *A. angustifrons* males: $\chi^2 = 47.78$, $df = 2$, $P < 0.001$, *A. sp. near pseudococci* females: $\chi^2 = 89.38$, $df = 2$, $P < 0.001$, *A. sp. near pseudococci* males: $\chi^2 = 62.02$, $df = 2$, $P < 0.001$) (Fig. 3). Females and males lived longer when provided with sucrose than with honeydew or water. *Delottococcus aberiae* honeydew did not increase the longevity of *A. angustifrons* females and males and only marginally increased the longevity of *A. sp. near pseudococci* females and males (between one and two days) when compared with water (Table 2).

DISCUSSION

The three generalist parasitoids, *A. angustifrons*, *A. sp. near pseudococci* and *L. algerica*, detected and accepted the new citrus pest *D. aberiae* but both behaviors varied among parasitoids and depended also on the mealybug instar. Among the three parasitoid species, *A. sp. near pseudococci* detected and accepted *D. aberiae* at higher rates, especially the third instar. This generalist parasitoid tends to accept mealybugs independently of their geographical origin, phylogenetic relationships and defensive behaviors (Bugila et al. 2015). *Leptomastix algerica* also tended to accept *D. aberiae* nymphs but it did not detect them easily. Unfortunately, the eggs of both parasitoids are encapsulated by *D. aberiae* (Tena et al. 2017). On the other hand, *A. angustifrons* females did neither detect nor accept *D. aberiae* nymphs and only accepted 40% of

the adults. This parasitoid has been observed on mealybugs from genus *Dysmicoccus* (Trjapitzin 2008) and it might not be adapted to parasitize pseudococcids from other genera. Although its long oviposition time and defensive responses of *D. aberiae* were not an impediment to successfully parasitize the adults, *A. angustifrons* does seem be able to detect *D. aberiae* and, moreover, its eggs are also encapsulated by *D. aberiae* (Tena et al. 2017).

Delottococcus aberiae responded to parasitoid attacks with four non-excluding active defensive behaviors: wriggling, swiveling around the inserted stylet, withdrawing the stylet and running away and secreting defensive fluids. These defensive behaviors had previously been described in other mealybug species (Boavida et al. 1995, Bokonon-Ganta et al. 1995, Cadée and van Alphen 1997, Bugila et al. 2014, Beltrà et al. 2015a). However, these defensive responses did not allow *D. aberiae* to escape parasitism from any parasitoid. Even *A. angustifrons* females, which needed more than 15 minutes to parasitize, were not disturbed by *D. aberiae* active defenses. Moreover, and contrary to previous studies with other mealybug species (Cadée and van Alphen 1997, Beltrà et al. 2015a), we did not find differences between mealybug instars. With all this said, the active defensive behavior of *D. aberiae* described in this paper cannot explain the low degrees of parasitism found in Spanish citrus since the detection of the mealybug in 2009 (Tena et al. 2014). Furthermore, our results support the hypothesis that most active host defenses might have not evolved as a response to parasitoid selective pressure but to other biological functions that also provide some protection against parasitoids (Gross 1993, Bugila et al. 2014). If there were specific defensive behaviors, we would expect differences among the responses to the three parasitoids.

Honeydew excreted by *D. aberiae* is a poor carbohydrate source for the studied parasitoids. Parasitoids fed honeydew extended their life for just one day more than unfed parasitoids and their lifespan was much shorter than those fed sucrose 2M (between 25 and 35 days long). The low nutritional suitability of honeydew may be based on plant-derived compounds (primary and secondary metabolites) as well as compounds synthesized by honeydew-producers (Wäckers et al. 2008). It is important to highlight that another mealybug *P. citri* excretes honeydew of high

value for parasitoids and ants in citrus (Pekas et al. 2011, Tena et al. 2013, Tena et al. 2016). Therefore, either *D. aberiae* or the rearing conditions are responsible for the poor honeydew. In our study, we have tested honeydew excreted by *D. aberiae* reared on detached fruits that could be of poorer quality for the mealybug than growing fruits in the field. In this scenario, it is reasonable to believe that honeydew composition may be affected. Our field observations, however, also show that ants do not attend *D. aberiae* colonies as frequently as they attend *P. citri* (A.T. personal observations), suggesting that *D. aberiae* excretes a poor quality honeydew also in the field. This is because ants tend hemipterans that excrete honeydew of high quality for parasitoids (Tena et al. 2016). Further research in the field is necessary to corroborate this result because sugar sources are crucial for the primary parasitoids of mealybugs, which are generally synovigenic. For example, *A. sp. near pseudococci* is synovigenic, emerge without mature eggs, and females lay few eggs per day (~5-7 per day) but for an extended period of time (~40 days) (Tingle and Copland, 1989).

Finally, we did not observe host-feeding in any parasitoid species. However, we cannot exclude the possibility of host-feeding by *A. sp. nr. pseudococci* and *L. algerica* in younger host instars, such as first and second instars, as our observations were carried out only on third instar nymphs and pre-reproductive adult females. For example, Karamaouna and Copland (2000) and Bokonon et al. (1995) observed females of *Leptomastix epona* (Walker) and *Anagyrus mangicola* (Noyes) (Hymenoptera: Encyrtidae) feeding on young instars of mealybugs in which they do not oviposit. If this is the case, host-feeding by *A. sp. near pseudococci* might cause the mortality of *D. aberiae* nymphs in Spanish citrus, where the parasitoid develops on *P. citri* (Martínez-Ferrer 2003, Campos-Rivela 2008), which coexist with *D. aberiae*. The importance of non-reproducing effects of parasitoids such as host feeding and mutilation or death of host probed and rejected by parasitoids are an underappreciated component of biological control services provided by parasitoids (Abram et al. 2016, Cebolla et al. 2017). The impact of host-feeding and other non-reproducing effects of *A. sp. near pseudococci* on *D. aberiae* should be

studied under field and laboratory conditions to unravel the effect of this abundant parasitoid in Europe.

ACKNOWLEDGMENTS

We want to thank two anonymous referees for their detailed review of the manuscript. We also thank J. Catalán and P. Bru (IVIA) for technical assistance and Dr. J. Calvo from Koppert BS for providing the *A. acerophagus*, *A. sp. near pseudococci* and *L. algerica* individuals. This research was partially funded by an INIA project (Project No. RTA2014-00067) and the Conselleria d'Agricultura, Pesca i Alimentació de la Generalitat Valenciana.

REFERENCES

- Abram, P. K., J. Brodeur, V. Burte, and G. Boivin. 2016.** Parasitoid-induced host egg abortion: An underappreciated component of biological control services provided by egg parasitoids. *Biol. Control* 98: 52-60.
- Avidov, Z., M. Balshin, and U. Gerson. 1970.** Studies on *Aphytis coheni*, a parasite of the California red scale, *Aonidiella aurantii*, in Israel. *Entomophaga* 15: 191-207.
- Bartlett, B.R. 1978.** Pseudococcidae. pp. 137-171. In Clausen, C.P. (ed), Introduced parasites and predators of arthropod pest and weeds: a world review. U. S. Department of Agriculture, Handbook No. 480, Washington, D.C. USA.
- Beltrà, A., F. Garcia-Marí, and A. Soto. 2013.** El cotonet de Les Valls, *Delottococcus aberiae*, nueva plaga de los cítricos. *Levante Agrícola* 419: 348-352.
- Beltrà, A., A. Soto, and A. Tena. 2015a.** How a slow-ovipositing parasitoid can succeed as a biological control agent of the invasive mealybug *Phenacoccus peruvianus*: implications for future classical and conservation biological control programs. *BioControl* 60: 473-484.
- Beltrà, A., P. Addison, J. A. Ávalos, D. Crochard, F. Garcia-Marí, E. Guerrieri, J.H. Giliomee, T. Malausa, C. Navarro-Campos, F. Palero, and A. Soto. 2015b.** Guiding Classical Biological Control of an Invasive Mealybug Using Integrative Taxonomy. *PLoS ONE* 10: e0128685. doi:10.1371/journal.pone.0128685.
- Blumberg, D. 1997.** Parasitoid encapsulation as a defense mechanism in the Coccoidea (Homoptera) and its importance in biological control. *Biol. Control* 8: 225-236.
- Blumberg, D., and R.G. Van Driesche. 2001.** Encapsulation rates of three encyrtid parasitoids by three mealybug species (Homoptera: Pseudococcidae) found commonly as pests in commercial greenhouses. *Biol. Control* 22: 191-199.
- Boavida, C., Ahounou, M., Vos, M., Neuenschwander, P., and J.J.M. van Alphen. 1995.** Host stage selection and sex allocation by *Gyransoidea tebygi* (Hymenoptera: Encyrtidae), a parasitoid of the mango mealybug, *Rastrococcus invadens* (Homoptera: Pseudococcidae). *Biol. Control* 5: 487-496.

- Bokonon-Ganta, A.H., P. Neuenschwander, J.J.M. van Alphen, and M. Vos. 1995.** Host stage selection and sex allocation by *Anagyrus mangicola* (Hymenoptera: Encyrtidae), a parasitoid of the mango mealybug, *Rastrococcus invadens* (Homoptera: Pseudococcidae). *Biol. Control* 5: 479–486.
- Bugila, A.A., J.C. Franco, E.B. da Silva, and M. Branco. 2014.** Defense response of native and alien mealybugs (Hemiptera: Pseudococcidae) against the solitary parasitoid *Anagyrus* sp. nr. *pseudococci* (Girault) (Hymenoptera: Encyrtidae). *J. Insect Behav.* 27: 439–453.
- Bugila, A.A., J.C. Franco, E.B. da Silva, and M. Branco. 2015.** Suitability of five mealybug species (Hemiptera: Pseudococcidae) as hosts for the solitary parasitoid *Anagyrus* sp. nr. *pseudococci* (Girault) (Hymenoptera: Encyrtidae). *Biocontrol Sci. Technol.* 25: 108–120.
- Bynum, E.K. 1937.** *Pseudococcobius terryi* Fullaway, a Hawaiian parasite of gray sugarcane mealybug in the United States. *J. Econ. Entomol.* 30: 756–761.
- Cadée, N., and J.J.M. van Alphen. 1997.** Host selection and sex allocation in *Leptomastidea abnormis*, a parasitoid of the citrus mealybug *Planococcus citri*. *Entomol. Exp. Appl.* 83: 277–284.
- Campos-Rivela, J. M. 2008.** Estudios comparativos de algunos aspectos de la biología de dos parasitoides: *Anagyrus pseudococci* (Girault) y *Leptomastix dactylopii* Howard (Hymenoptera: Encyrtidae), para la mejora de los cítricos *Planococcus citri* Risso (Hemiptera: Pseudococcidae). PhD dissertation, Universitat Jaume I, Castellón, Spain.
- Cebolla, R., P. Vanaclocha, A. Urbaneja, and A. Tena. 2017.** Overstinging by hymenopteran parasitoids causes mutilation and surplus killing of hosts. *J. Pest Sci.* DOI: 10.1007/s10340-017-0901-9.
- Crawley, M.J. 2007.** *The R Book*. John Wiley & Sons, New York. USA.
- Daane, K.M., K.R. Sime, J. Fallon, and M.L. Cooper. 2007.** Impacts of Argentine ants on mealybugs and their natural enemies in California's coastal vineyards. *Ecol. Entomol.* 32: 583–596.
- De Farias, A.M., and Hopper K.R. 1999.** Oviposition behavior of *Aphelinus asychis* (Hymenoptera: Aphelinidae) and *Aphidius matricariae* (Hymenoptera: Aphidiidae) and defense behavior of their host *Diuraphis noxia* (Homoptera: Aphididae). *Environ. Entomol.* 28: 858–862.
- Desneux, N., R. J. Barta, K. A. Hoelmer, K. R. Hopper, and G. E. Heimpel. 2009.** Multifaceted determinants of host specificity in an aphid parasitoid. *Oecologia* 160: 387–398.
- Foldi, I. 1997.** Defense strategies in scale insects: phylogenetic inference and evolutionary scenarios (Hemiptera, Coccoidea). pp 203–230. In Grandcolas, P. (ed) *The origin of biodiversity in insects: phylogenetic tests of evolutionary scenarios*, 1st edn. Museum National d'Histoire Naturelle, Paris, France.
- Franco, J.C., P. Suma, E.B. Da Silva, D. Blumberg, and Z. Mendel. 2004.** Management strategies of mealybug pests of citrus in Mediterranean countries. *Phytoparasitica* 32: 507–522.
- Gross, P. 1993.** Insect behavioral and morphological defenses against parasitoids. *Annu. Rev. Entomol.* 38: 251–273.

386 **Gullan, P.J., and M. Kosztarab .1997.** Adaptations in scale insects. *Annu. Rev. Entomol.*
387 42:23–50.

388 **Gutierrez, A.P., K.M. Daane, L. Ponti, V.M. Walton, and C.K. Ellis. 2008.** Prospective
389 evaluation of the biological control of vine mealybug: refuge effects and climate. *J. Appl.*
390 *Ecol.* 45: 524-536.

391 **Hogervorst, P.A.M., F.L. Wäckers, and J. Romeis. 2007.** Detecting nutritional state and food
392 source use in field-collected insects that synthesize honeydew oligosaccharides. *Funct.*
393 *Ecol.* 21: 936–946.

394 **Islam, K.S., and M.J.W. Copland. 1997.** Host preference and progeny sex ratio in a solitary
395 koinobiont mealybug endoparasitoid, *Anagyrus pseudococci* (Girault), in response to its
396 host stage. *Biocontrol Sci. Technol.* 7: 449-456.

397 **Karamaouna, F., and M.J.W. Copland. 2000.** Oviposition behavior, influence of experience
398 on host size selection, and niche overlap of the solitary *Leptomastix epona* and the
399 gregarious *Pseudaphycus flavidulus*, two endoparasitoids of the mealybug *Pseudococcus*
400 *viburni*. *Entomol. Exp. Appl.* 97: 301-308.

401 **Martínez-Blay, V., J. Pérez-Rodríguez, A. Tena, and A. Soto. 2017.** Density and phenology
402 of the invasive mealybug *Delottococcus aberiae* De Lotto (Hemiptera: Pseudococcidae) on
403 citrus: implications for its integrated pest management. Accepted in *J. Pest Sci.*

404 **Martínez-Ferrer, M.T. 2003.** Biología y control del cotonet *Planococcus citri* (Homoptera:
405 Pseudococcidae) en huertos de cítricos. PhD dissertation, Universitat Politècnica de
406 València, Valencia, Spain.

407 **Miller, D.R., and J.H. Giliomee. 2011.** Systematic revision of the mealybug genus
408 *Delottococcus* Cox & Ben-Dov. *Afr. Entomol.* 19: 614-640.

409 **Moore, D. 1988.** Agents used for biological control of mealybugs (Pseudococcidae). *Biocontrol*
410 *News Info* 9: 209-225.

411 **Muniappan, R., D.E. Meyerdirk, F.M. Sengebau, D.D. Berringer, and G.V.P. Reddy. 2006.**
412 Classical biological control of the papaya mealybug, *Paracoccus marginatus* (Hemiptera:
413 Pseudococcidae) in the Republic of Palau. *Fla. Entomol.* 89: 212–217.

414 **Noyes, J. S., and M. Hayat. 1994.** Oriental mealybug parasitoids of the Anagyrini
415 (Hymenoptera: Encyrtidae). Cab International, Wallingford. UK.

416 **Pekas, A., A. Tena, A. Aguilar, and F. Garcia-Marí. 2011.** Spatio-temporal patterns and
417 interactions with honeydew-producing Hemiptera of ants in a Mediterranean citrus
418 orchard. *Agric. For. Entomol.* 13: 89–97.

419 **Pérez-Rodríguez, J., V. Martínez-Blay, A. Soto, J. Selfa, C. Monzó, A. Urbaneja, and A.**
420 **Tena. 2017.** Aggregation patterns, sampling plan, and economic injury levels for the new
421 citrus pest *Delottococcus aberiae* De Lotto. *J. Econ. Entomol.* DOI: [10.1093/jee/tox258](https://doi.org/10.1093/jee/tox258).

422 **Sime, K.R., and K.M. Daane. 2014.** Rapid, non-discriminatory oviposition behaviors are
423 favored in mealybug parasitoids when Argentine ants are present. *Environ. Entomol.* 43:
424 995–1002.

425 **Tena, A., A. Beltrà, and A. Soto. 2012.** Novel defenses of *Protopulvinaria pyrifomis*
426 (Hemiptera: Coccidea) against its major parasitoid *Metaphycus helvolus* (Hymenoptera:
427 Encyrtidae): Implications for biological control of soft scales. *Biol. Control* 62: 45-52.

428 **Tena, A., A. Pekas, F.L. Wäckers, and A. Urbaneja. 2013.** Energy reserves of parasitoids
429 depend on honeydew from non-hosts. *Ecol. Entomol.* 38: 278-289.

430 **Tena, A., J. Catalán, P. Bru, and A. Urbaneja. 2014.** *Delottococcus aberiae*, nueva plaga de
431 cítricos. *Agricultura 978*: 746-748.

432 **Tena, A., F.L. Wäckers, G.E. Heimpel, A. Urbaneja, and A. Pekas. 2016.** Parasitoid
433 nutritional ecology in a community context: the importance of honeydew and implications
434 for biological control. *Curr. Opin. Insect Sci.* 14: 100-104.

435 **Tena, A., J. García-Bellón, and A. Urbaneja. 2017.** Native and naturalized mealybug
436 parasitoids fail to control the new citrus mealybug pest *Delottococcus aberiae*. *J. Pest Sci.*
437 90: 659-667.

438 **Tingle, C.C., and M.J. Copland. 1989.** Progeny production and adult longevity of the
439 mealybug parasitoids *Anagyrus pseudococci*, *Leptomastix dactylopii*, and *Leptomastidea*
440 *abnormis* [Hym.: Encyrtidae] in relation to temperature. *BioControl* 34: 111-120.

441 **Trjapitzin, V.A. 2008.** A review of encyrtid wasps (Hymenoptera, Chalcidoidea, Encyrtidae) of
442 Macaronesia. *Entomologicheskoe Obozrenie* 87: 166-184

443 **van Driesche, R.G., A. Belloti, C.J. Herrera, and J.A. Castello .1987.** Host feeding and
444 ovipositor insertion as sources of mortality in the mealybug *Phenacoccus herreni* caused
445 by two encyrtids *Epidinocarsis diversicornis* and *Acerophagus coccois*. *Entomol. Exp.*
446 *Appl.* 44: 97–100.

447 **Vanaclocha, P., S. Vacas, C. Alfaro, J. Primo, M.J. Verdú, V. Navarro-Llopis, and A.**
448 **Urbaneja. 2012.** Life history parameters and scale-cover surface area of *Aonidiella*
449 *aurantii* are altered in a mating disruption environment: implications for biological control.
450 *Pest Manag. Sci.* 68: 1092-1097.

451 **Vet, L.E., and M. Dicke. 1992.** Ecology of infochemical use by natural enemies in a tritrophic
452 context. *Annu. Rev. Entomol.* 37: 141–172.

453 **Wäckers, F.L., P.C.J. van Rijn, and G.E. Heimpel. 2008.** Honeydew as a food source for
454 natural enemies: making the best of a bad meal? *Biol. Control* 45: 176-184.

455 **Wyckhuys, K.A.G., L. Stone, N. Desneux, K.A. Hoelmer, K.R. Hopper, and G.E. Heimpel.**
456 **2008.** Parasitism of the soybean aphid *Aphis glycines* by *Binodoxys communis*: the role of
457 aphid defensive behavior and parasitoid reproductive performance. *Bull. Entomol. Res.* 98:
458 361–370.

459

Figure legends

Figure 1. Mean time spent ($\pm$ SE) by *Acerophagus angustifrons*, *Anagyrus* sp. near *pseudococci* and *Leptomastix algerica* when they parasitized 3rd instar nymphs (N3) and preovipositing adult females (F1) of the mealybug *Delottococcus aberiae*.

Fig. 2. Mean number of defensive responses ($\pm$ SE) of 3rd instar nymphs (N3) and preovipositing adult females (F1) of the mealybug *Delottococcus aberiae* when attacked by the parasitoid species *Acerophagus angustifrons*, *Anagyrus* sp. near *pseudococci* and *Leptomastix algerica*.

Fig. 3. Survival of females and males of the parasitoids *Acerophagus angustifrons* and *Anagyrus* sp. near *pseudococci* fed on sucrose 2M (mol/L), honeydew excreted by *Delottococcus aberiae* or water.

Table 1. Proportion of *Delottococcus aberiae* ($\pm$ SE) detected and accepted by the parasitoids *Acerophagus angustifrons*, *Anagyrus* sp. near *pseudococci* and *Leptomastix algerica* when encountering 3rd instar nymphs (N3) and preovipositing adult females of the mealybug (F1). Different letters within a host instar indicates significant differences among parasitoid species ($P < 0.05$).

Host instar	Parasitoid species	Detection	Acceptance
N3	<i>A. angustifrons</i>	0.23 $\pm$ 0.12 ^b	0.08 $\pm$ 0.08 ^b
	<i>A. sp. near pseudococci</i>	0.94 $\pm$ 0.06 ^a	0.78 $\pm$ 0.10 ^a
	<i>L. algerica</i>	0.55 $\pm$ 0.11 ^b	0.55 $\pm$ 0.11 ^a
F1	<i>A. angustifrons</i>	0.67 $\pm$ 0.14 ^a	0.42 $\pm$ 0.15 ^b
	<i>A. sp. near pseudococci</i>	0.88 $\pm$ 0.08 ^a	0.82 $\pm$ 0.10 ^a
	<i>L. algerica</i>	0.67 $\pm$ 0.10 ^a	0.67 $\pm$ 0.10 ^{ab}

Table 2. Mean longevity in days ($\pm$ SE) of females and males of the parasitoids *Acerophagus angustifrons* and *Anagyrus* sp. near *pseudococci* when fed on sucrose 2M, honeydew excreted by *Delottococcus aberiae* and water. Different letters within the same row denote significant differences between diets ($P < 0.05$).

Especies	Sex	Diet		
		Sucrose	Honeydew	Water
<i>A. angustifrons</i>	Female	30.8 $\pm$ 1.9 ^a	2.3 $\pm$ 0.18 ^b	1.8 $\pm$ 0.18 ^b
	Male	26.9 $\pm$ 1.45 ^a	1.7 $\pm$ 0.17 ^b	1.5 $\pm$ 0.14 ^b
<i>A. sp. near pseudococci</i>	Female	46.0 $\pm$ 3.52 ^a	4.2 $\pm$ 0.48 ^b	2.3 $\pm$ 0.16 ^c
	Male	28.7 $\pm$ 2.39 ^a	3.7 $\pm$ 0.49 ^b	1.9 $\pm$ 0.23 ^c