

ORIGINAL RESEARCH

Influence of farm management on the dynamics of *Salmonella enterica* serovar Infantis shedding and antibiotic resistance during the growing period of broiler chickens

Laura Montoro-Dasi^{1,2}  | Arantxa Villagra³ | Santiago Vega⁴ | Clara Marin⁴

¹ Instituto de Ciencia y Tecnología Animal, Universidad Politécnica de Valencia, Valencia, Spain

² Centro de Calidad Avícola y Alimentación Animal de la Comunidad Valenciana (CECAV), Castellón, Spain

³ Centro de Investigación y Tecnología Animal, Instituto Valenciano de Investigaciones Agrarias, Castellón, Spain

⁴ Departamento de Producción y Sanidad Animal, Salud Pública Veterinaria y Ciencia y Tecnología de los Alimentos, Instituto de Ciencias Biomédicas, Facultad de Veterinaria, Universidad Cardenal Herrera-CEU, CEU Universities, Moncada, España

Correspondence

Clara Marin Orenge, Departamento de Producción y Sanidad Animal, Salud Pública Veterinaria y Ciencia y Tecnología de los Alimentos, Instituto de Ciencias Biomédicas, Facultad de Veterinaria, Universidad Cardenal Herrera-CEU, CEU Universities, Avenida Seminario s/n, Moncada, 46113, España. Email: clara.marin@uchceu.es

Funding information

National Institute for Agricultural Research and Experimentation; Ministry of Economy, Industry and Competitiveness, Grant/Award Number: RTA2017-00013; University CEU-UCH, Grant/Award Number: INDI19/32

[The copyright line for this article was changed on 21 June 2021 after original online publication.]

INTRODUCTION

Animal welfare and food safety are increasing concerns for poultry product consumers.¹ Both issues are closely related, as it has been demonstrated that if animals are in good welfare status, their resilience is increased, and they can cope with environmental challenges or infectious diseases.^{2–5} For that reason, an investment in more efficient and animal friendly management measures in the poultry sector could directly affect animal health.^{3,6–9} In this sense, a good ventilation system is essential for heat stress manage-

Abstract

Background: *Salmonella enterica* serovar Infantis is a zoonotic pathogen isolated in broilers causing great economic losses in the European poultry sector. It is demonstrated that an investment in management measures at farm level could directly affect the control of food chain microorganisms. The aim of this study was to investigate the development of *S. Infantis* antimicrobial resistance (AMR) patterns during the growing period, according to flock density and ventilation management, without antibiotic administration.

Methods: The experiment was performed in two identical poultry houses, evaluating commercial and optimal farm conditions. At 24 h of rearing, 20% of the animals were orally infected with a *S. Infantis* strain susceptible to all the antibiotics tested. To study *Salmonella* shedding, faeces samples from each experimental group were taken weekly and analysed as per ISO/TS 6579-2:2017. Antibiotic susceptibility was assessed according to Decision 2013/653.

Results: *Salmonella* shedding showed that the lowest counts were observed in the first week post-infection and highest at slaughter day for both groups. Moreover, 100% of the isolates were multi-resistant.

Conclusion: The acquisition of AMR by *S. Infantis* starts at the onset of the production cycle and is maintained until the end, demonstrating the importance of transmission of AMR in zoonotic bacteria at farm level.

KEYWORDS

antimicrobial resistance, broiler, growing period, multidrug-resistance, salmonella

ment, a factor that undermines the productivity and immunology of livestock.^{10,11} Likewise, high stocking density also has an adverse effect on the performance and immune status of broilers.^{12–14}

Furthermore, despite the strict legislation against *Salmonella*, these bacteria remain the principal source of human foodborne disease in Europe, and poultry products are the main source involved in human outbreaks.^{1,15,16} Moreover, *Salmonella enterica* serovar Infantis is an emerging serovar of great concern for European broiler production, as it has been demonstrated that this serovar is present in 50%

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Veterinary Record* published by John Wiley & Sons Ltd on behalf of British Veterinary Association

of *Salmonella* contaminated broiler meat samples analysed.^{16,17} Consequently, nowadays *S. Infantis* control at farm level is one of the main objectives for the poultry sector.

One hypothesis that explains the emergence of *S. Infantis* in the poultry sector is its ability to gain antimicrobial resistance (AMR) from the gut microbiota and/or environment.^{18–20} Moreover, a novel megaplasmid has been identified that represents a recent evolutionary change in the pathogenicity and stress tolerance of local *S. Infantis* population.²¹ In this context, AMR control in the field requires effective surveillance programs, proper food handling practices and prudent use of antibiotics throughout the production cycle.^{22,23} However, to be able to establish adequate control measures, it is necessary to have better knowledge of the epidemiology of this serovar.

In accordance with the increasing consumer concern for animal welfare and the public health issue of AMR, the objective of this study was to investigate the development of *S. Infantis* AMR during the broiler growing period, according to density and ventilation management.

METHODS

In this experiment, all animals were handled according to the principles of animal care published by Spanish Royal Decree 53/2013.²⁴

Experiment design

The study was performed in two identical poultry houses of an experimental poultry farm at the Center for Research and Animal Technology (CITA-IVIA, in its Spanish acronym *Valencian Institute for Agrarian Research*, Segorbe, Spain). For this purpose, two different environmental farm conditions, commercial farm conditions (CFC) (house 1) and optimal farm conditions (OFC) (house 2), were evaluated. For CFC, chicks were housed at 35 kg/m² density, and non-optimal parameters of ventilation were applied (allowing a maximum concentration of ammonia of 25 ppm). While in OFC, the animals were housed in low density at 17 kg/m², and ventilation was provided within the optimal parameters (allowing a maximum concentration of ammonia of 10 ppm).

To this end, day-old-chicks (Ross) (males and females) were distributed in two identical poultry houses ($n = 1062$, 531 per house). Within each of the houses, 204 of 531 animals were located in 12 pens with wood shavings as bedding material. The rest of the animals (327/531) were housed in the remaining space using also wood shavings as bedding material to simulate production conditions. The house was supplied with programmable electrical lights, automated electric heating and forced ventilation. The environmental temperature was gradually decreased from 32°C (1 day) to 19°C (42 days) following common practice in poultry production. The experimental pelleted feed was commercial feed according to standard diets for broilers. Two different diets were

offered to the birds: starter (1–21 days) and grower (21–42 days). Only one batch of feed per age (starter and grower) was provided. Nutritional and product analysis were assessed before the arrival of animals. Feed was weighed, manually distributed and added *ad libitum*. Furthermore, the mortality and the presence of diarrhea were registered daily. Finally, animals were weighed at weekly intervals, and feed consumption per pen was recorded.

Salmonella infection

At 24 h after placing, 20% of birds/pen were orally infected with *S. Infantis*. The experimental infection was done with 100 µl of a *S. Infantis* diluted at an infective titer of 10⁴ CFU/ml. The strain was selected from a database of *Salmonella* strains isolated from the *Salmonella* National Control Program (CECAV, in its Spanish acronym *Centro de Calidad Avícola y Alimentación Animal de la Comunidad Valenciana*, Castellón, Spain). To ensure that this strain was susceptible to all antibiotics studied, antimicrobial susceptibility was tested according to the European Committee on Antimicrobial Susceptibility Testing guidelines.²⁵ The source for zone diameters used for interpretation of the test was http://www.eucast.org/clinical_breakpoints/. The strain of *S. Infantis* was inoculated into Mueller-Hinton agar (Scharlab, S.L., Barcelona, Spain) to form a bacterial lawn, the antibiotic disks were added, and plates were incubated at 37°C ± 1°C for 24 ± 3 h.

Salmonella detection and identification

Salmonella status of the chicken houses was tested before the arrival of the animals in accordance with ISO 6579-1:2017.²⁶ In addition, *Salmonella* status of the flock was tested at the arrival day, collecting samples of meconium ($n = 250$) and delivery box liners ($n = 10$).²⁷

Salmonella enumeration was assessed as per ISO/TS 6579-2:2017.²⁸ Animals were sampled at different times throughout the growing period (7, 14, 21, 28, 35 and 42 days of age). For each sampling time and house (CFC vs. OFC), faeces samples (25 g) were directly collected from each pen per duplicate ($n = 24$). Once in the laboratory, two pools of samples from each replicate per house ($n = 2$ pools/treatment/house) were homogenized and transferred into 225 ml of Buffered Peptone Water (Scharlab, S.L., Barcelona, Spain). Afterwards, 2.5 ml of the suspension was transferred into an empty tube. Serial 1:5 dilutions were made from each tube and incubated at 37°C for 18 ± 2 h. After incubation, 20 µl was transferred onto Rappaport Vasiliadis agar plates (MSRV, Difco, Valencia, Spain) and incubated at 41.5°C for 24–48 h. Suspected plates were streaked into XLD medium (Scharlab, S.L., Barcelona, Spain) and incubated at 37°C ± 1°C for 24 ± 3 h. Then, API-20E test (Biomerieux, S.L., Barcelona, Spain) was performed to confirm *Salmonella*. Finally, for the estimation of Most Probable Number, the software described by




FIGURE 1 *Salmonella* excretion dynamic in CFC and OFC during growing period. ^{a,b}: different superscripts means significant differences with a p -value < 0.05

Jarvis et al was used, and the results were transformed into logarithms (log₁₀ CFU/g).²⁹ To confirm that the isolates were obtained from the original inoculum, *Salmonella* strains were serotyped at CECAV, using the Kauffman-White-Le Minor scheme.³⁰

Antimicrobial susceptibility test

Antimicrobial susceptibility of the strains isolated was tested as reported above. The antibiotics selected were those set forth in Decision 2013/652³¹ including two quinolones: ciprofloxacin (CIP) (5 µg) and nalidixic acid (NAL) (30 µg); three b-lactams: ampicillin (AMP) (10 µg), cefotaxime (CTX) (30 µg) and ceftazidime (CAZ) (30 µg); one phenicol: chloramphenicol (CHL) (5 µg); one potentiated sulfonamide: trimethoprim-sulfamethoxazole (SXT) (1.25/23.75 µg); one polymyxin: colistin (CST) (10 µg); one macrolide: azithromycin (AZM) (15 µg); one glycylicline: tigecycline (TGC) (15 µg); one aminoglycoside: gentamycin (GEN) (10 µg) and one pyrimidine: trimethoprim (TMP) (5 µg). MDR was defined as acquired resistance to at least one agent in two or more antimicrobial classes.³²

Statistical analysis

ANOVA test was used to study the dynamics of *S. Infantis* shedding and AMR during growing period

under different farm conditions (CFC and OFC). A p -value < 0.05 was considered to indicate a statistically significant difference. Analyses were carried out using a commercially available software application (SPSS 24.0 software package; SPSS Inc., Chicago, IL, 2002).

RESULTS

During this experiment, all the productive parameters were according to the breed standards, and no signs of intestinal disease were observed. Thus, no antibiotics were administered in this study.

Salmonella excretion

At the start of the trial, negative *Salmonella* status of the chicken houses and the day-old-chickens was confirmed. Moreover, all the *Salmonella* strains isolated during this study were serotyped as *S. Infantis*.

Results obtained for CFC and OFC are presented in Figure 1. For both environmental farm conditions studied, the lowest excretion of *S. Infantis* was observed in the first week post-infection (wpi). Then, for CFC, *S. Infantis* detection increased until 14 days and then became stable until the end of growing period (p -value < 0.05). However, for OFC, *S. Infantis* counts increased until 21 days of the growing period and then remained stable

TABLE 1 Antibiotic resistance isolates according to the antibiotic and the moment of the growing period in commercial (CFC) and optimal farm conditions (OFC)

Environmental conditions	wpi	N pools	CIP	NAL	CTX	CAZ	AMP	CHL	SXT	CST	AZM	TGC	GEN	TMP
CFC	1	2	2	2	2	1	0	0	2	0	0	0	0	2
	2	2	2	2	1	1	0	0	2	1	0	0	0	2
	3	2	2	2	0	0	0	0	2	0	0	0	0	2
	4	2	2	2	0	1	0	0	2	0	0	0	0	2
	5	2	2	2	0	0	0	0	1	0	0	0	0	2
	6	2	2	2	0	0	0	0	2	0	2	0	0	2
	Total	12	12	12	3	3	0	0	11	1	2	0	0	0
OFC	1	2	2	2	2	1	0	0	2	0	0	0	0	2
	2	2	2	2	1	0	1	1	2	0	0	0	0	1
	3	2	2	2	0	0	0	0	2	0	0	0	0	2
	4	2	2	2	1	1	1	0	2	0	0	0	0	2
	5	2	2	2	0	0	0	0	2	0	0	0	0	2
	6	2	2	2	0	0	0	0	2	0	2	0	0	2
	Total	12	12	12	4	2	2	1	12	0	2	0	0	0

Abbreviations: AMP, ampicillin; AZM, azithromycin; CAZ, ceftazidime; CHL, chloramphenicol; CIP, ciprofloxacin; CST, colistin; CTX, cefotaxime; GEN, gentamycin; NAL, nalidixic acid; SXT, trimethoprim-sulfamethoxazole; TGC, tigecycline; TMP: trimethoprim.

until the end of growing (p -value < 0.05). However, no statistically significant differences were found between treatments (CFC vs. OFC) in *Salmonella* counts (p -value > 0.05).

Prevalence of antibiotic resistance and multidrug-resistance

Although the *S. Infantis* strain used to infect the animals was completely susceptible to all antibiotics tested at the time of infection, and no antibiotics were administered during the growing period, *Salmonella* isolates obtained from both groups ($n = 24$) were MDR after 1 wpi. No statistically differences were found between experimental conditions (CFC vs. OFC) and the time of sampling (p -value > 0.05).

For CFC, the highest percentages of AMR were found to CIP (100%, $n = 12$), NAL (100%, $n = 12$) and TMP (100%, $n = 12$), followed by SXT (91.7%, $n = 11$), CTX (25.0%, $n = 3$), CAZ (25.0%, $n = 3$), AZM (16.7%, $n = 2$) and finally, CST (8.3%, $n = 1$). No resistance was found against AMP, CHL, GEN and TGC. Regarding resistance dynamic through the entire growing period, at 1 wpi, *S. Infantis* strains showed resistance to CIP, NAL, CTX, CAZ, SXT and TMP. It is important to highlight that at 2 wpi, resistance to CST also appeared. However, from the third wpi onwards, only resistance to CIP, NAL, SXT and TMP remained until slaughter day (Table 1).

In the case of OFC, the highest AMR percentages were observed to CIP (100%, $n = 12$), NAL (100%, $n = 12$) and SXT (100%, $n = 12$), followed by TMP (91.7%, $n = 11$). The remaining antibiotics showed a lower AMR percentage: CTX (33.3%, $n = 4$), CAZ (16.7%, $n = 2$), AMP (16.7%, $n = 2$), AZM (16.7%, $n = 2$)

and CHL (8.3%, $n = 1$). Regarding the AMR dynamics during the growing period, at 1 wpi and at the slaughter day, *S. Infantis* strains were resistant to the same antibiotics of CFC isolated strains. However, in OFC no resistance to CST appeared during the growing period (Table 1).

Abbreviations: AMP, ampicillin; AZM, azithromycin; CAZ, ceftazidime; CHL, chloramphenicol; CIP, ciprofloxacin; CST, colistin; CTX, cefotaxime; GEN, gentamycin; NAL, nalidixic acid; TGC, tigecycline; TMP, trimethoprim; SXT, trimethoprim-sulfamethoxazole.

Antibiotic resistance patterns

A number of *Salmonella* strains isolated resistant to the different antibiotics tested according to different environmental farm conditions (CFC and OFC) are presented in Table 2.

Overall, 11 different resistance patterns were observed. The combination of CIP-NAL-SXT-TMP (37.5%, $n = 9$) was the pattern most frequently observed, followed by CIP-NAL-SXT-AZM-TMP (16.67%, $n = 4$).

DISCUSSION

The present study examined the development of *S. Infantis* AMR in broiler chickens during the growing period, comparing two different environmental conditions according to density and ventilation parameters. To our best knowledge, this is the first study in scientific literature to evaluate the effect of these management measures at farm level on *S. Infantis* epidemiology.

TABLE 2 Number of *Salmonella* strains isolated resistant to the different antibiotics tested according to different environmental farm conditions

Environmental conditions	Number of AMR to the indicated number of antibiotics											Total
	0	1	2	3	4	5	6	7	8	9	10	
CFC	0	0	0	1	4	5	2	0	0	0	0	12
OFC	0	0	0	1	5	3	1	2	0	0	0	12

Abbreviations: AMR, antimicrobial resistance; CFC, commercial farm conditions; OFC, optimal farm conditions.

On the day of placement, the negative *Salmonella* status of the chickens was confirmed. After infection, for CFC experimental group *S. Infantis* counts increased until 14 days and, for OFC until 21 days, without statistically significant differences between treatments. Our results agree with those reported previously by Marin and Lainez, when *Salmonella* detection in faeces increased until second week of age, coinciding with the maturation of the animals' immune system and remaining stable until processing day.^{33,34}

In the European Union, a strict poultry welfare legislation has been set out at farm level.³⁵ However, a large section of society calls for a continuous increase in animal welfare during the grow-out period.¹ In fact, different authors indicate that lower stress situations increase the potential of the immune system to protect the individual against pathogens.^{3,4,14,17,36–39} However, the results of our study showed that the improvement in ventilation or density parameters of the flock has no effects in terms of either *Salmonella* shedding which is in line with Velasquez et al and Pulido-Landínez.^{40,41}

AMR rates of *Salmonella* isolates obtained since the start of the trial showed that no statistical differences were found between treatments, despite the improvement in management conditions. In addition, it is important to underscore that a high percentage of *S. Infantis* isolated during the growing period were MDR, although no antibiotics were administered.^{16,23,42}

Different hypotheses could explain this fact. Previous studies using genomic analysis of bacteria indicated they could acquire resistance profiles by incorporating different genetic elements through horizontal gene transfer from other bacteria and/or from the environment.^{34,43–48} In this sense, the commensal microbiota could acquire the AMR and, intestinal zoonotic bacteria such as *Salmonella* could acquire the AMR by conjugation, transformation or transduction mechanisms.^{49,50} For that reason, different scientific studies underline the importance of developing sanitary measures at the interface between the environment and livestock farming.^{51–53} However, further studies are needed to confirm the main source of AMR of the *Salmonella* strains at farm level.

Moreover, in reference to AMR percentages obtained from different antibiotics assessed, it is important to highlight the results obtained against CST and TGC, as they are considered critically important antimicrobials used as last-resort drugs to treat human infectious diseases.^{50,54} On the one hand, no iso-

lates showed AMR to TGC. The results agree with that reported by the European Food Safety Authority (EFSA),⁵⁰ and it might be explained by the restricted use to human in hospital treatments.⁵⁵ Conversely, the presence of AMR against CST could be due to its use in animal production for several years to treat infectious diseases and as a growth promotor⁵⁶ and, as indicated by previous studies, resistant genes could remain in the environment and reach the microbiota of animals, and from there transmitted to zoonotic bacteria. Furthermore, it is important to note that the highest AMR obtained is to CIP, NAL, SXT and TMP. In 2020, EFSA reported very high levels of resistance to CIP, NAL and SXT in *Salmonella* isolated from broilers, and low levels of resistance to AMP and CHL, matching our results.⁵⁰ Moreover, specifically for SXT and TMP, one hypothesis that could explain the results obtained in this study is that these antibiotics are permitted in Spain as therapeutic agents for antibacterial therapy in animal.⁵⁵ This study reveals the importance of AMR monitoring in zoonotic and commensal bacteria in food-producing animals and their food products to be able to understand the development and diffusion of resistance, providing relevant risk assessment data, and evaluating targeted interventions.^{50,57}

In conclusion, the results of this study showed that when chicks are infected with the serovar *S. Infantis* at day one of the growing period, they continue shedding the bacteria in faeces until the processing day. Besides, the acquisition of AMR began at the onset of the production cycle and continued until the end, regardless of different management conditions applied. Nevertheless, it is important to highlight that no molecular studies of the microbiota interaction have been done in this study, which may restrict the interpretation of the results obtained. Thus, further deeply studies of the plasmids, pathogenicity islands or transposons are needed to achieve a better knowledge of *S. Infantis* AMR dynamics at the farm level, in order to establish better control programs and reduce its prevalence throughout the food chain.

ACKNOWLEDGEMENTS

We wish to thank the 'Improvement of Production System-related Food Safety and End Products' research group (Veterinary Faculty, University CEU-Cardenal Herrera) for their technical support. English text version was revised by N. Macowan English Language Service. This research was supported by the National Institute for Agricultural Research and Experimentation and the Ministry of Economy, Industry and

Competitiveness (RTA 2017-00013, Program: MINECO, Ministerio de Economía y Competitividad) and University CEU-UCH (INDI 19/32).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

Data curation: L. Montoro-Dasi, C. Marin and S. Vega. Formal analysis: L. Montoro-Dasi and C. Marin. Funding acquisition: A. Villagra. Investigation: L. Montoro-Dasi and C. Marin. Methodology: A. Villagra and C. Marin. Writing-original draft: L. Montoro-Dasi and C. Marin. Writing-review and editing: A. Villagra, S. Vega and C. Marin. All authors have read and agreed to the published version of the manuscript.

ORCID

Laura Montoro-Dasi  <https://orcid.org/0000-0002-3130-0933>

REFERENCES

- Bhaisare DB, Thyagarajan D, Churchil RR, Punniamurthy N. Bacterial pathogens in chicken meat: review. *Int J Life Sci Res.* 2014;2:2348–313.
- Soleimani AF, Zulkifli I, Hair-Bejo M. Epigenetic modification: possible approach to reduce *Salmonella enterica* serovar enteritidis susceptibility under stress conditions. *Avian Pathol.* 2012;41:351–4.
- Gomes AVS, Quinteiro-Filho WM, Ribeiro A. Overcrowding stress decreases macrophage activity and increases *Salmonella Enteritidis* invasion in broiler chickens. *Avian Pathol.* 2014;43:82–90.
- Dawkins MS. Animal welfare and efficient farming: is conflict inevitable? *Anim Prod Sci.* 2017;57:201–8.
- Swaggerty CL, Callaway TR, Kogut MH, Piva A, Grilli E. Modulation of the immune response to improve health and reduce foodborne pathogens in poultry. *Microorganisms.* 2019;7:1–10.
- Guardia S, Konsak B, Combes S. Effects of stocking density on the growth performance and digestive microbiota of broiler chickens. *Poult Sci.* 2011;90:1878–89.
- Franz E, van der Fels-Klerx HJ, Thissen J, van Asselt ED. Farm and slaughterhouse characteristics affecting the occurrence of *Salmonella* and *Campylobacter* in the broiler supply chain. *Poult Sci.* 2012;91:2376–81.
- Ben SN, Averós X, Estevez I. Technology and poultry welfare. *Animals.* 2016;6:1–21.
- Rouger A, Tresse O, Zagorec M. Bacterial Contaminants of Poultry Meat: Sources, Species, and Dynamics. *Microorganisms.* 2017;5:50.
- Farag MR, Alagawany M. Physiological alterations of poultry to the high environmental temperature. *J Therm Biol.* 2018;76:101–6.
- Ranjan A, Sinha R, Devi I, Rahim A, Tiwari S. Effect of Heat Stress on Poultry Production and their Management Approaches. *Int J Curr Microbiol Appl Sci.* 2019;8:1548–55.
- Qaid M, Albatshan H, Shafey T, Hussein E, Abudabos AM. Effect of stocking density on the performance and immunity of 1-to 14-d- old broiler chicks. *Rev Bras Cienc Avic.* 2016;18:683–92.
- Farhadi D, Hosseini SM, Dezfuli BT. Effect of house type on growth performance, litter quality and incidence of foot lesions in broiler chickens reared in varying stocking density. *J BioSci Biotech.* 2016;5:69–78.
- Desoky AA. Growth Performance and Immune Response of Broiler Chickens Reared Under High Stocking Density and Vitamin E Supplementation. *Egypt Poult Sci J.* 2018;38:607–20.
- EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). Opinion of the scientific panel on biological hazards (BIOHAZ) related to the use of vaccines for the control of *Salmonella* in poultry. *EFSA J.* 2004;2:1–74.
- EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union One Health 2018 Zoonoses Report. *EFSA J.* 2019;17:e05926.
- EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). *Salmonella* control in poultry flocks and its public health impact. *EFSA J.* 2019;17:e05596.
- Shah DH, Paul NC, Sischo WC, Crespo R, Guard J. Population dynamics and antimicrobial resistance of the most prevalent poultry-associated *Salmonella* serotypes. *Poult Sci.* 2016;96:687–702.
- Abdi RD, Mengstie F, Beyi AF, Beyene T, Waktole H, Mammo B, et al. Determination of the sources and antimicrobial resistance patterns of *Salmonella* isolated from the poultry industry in Southern Ethiopia. *BMC Infect Dis.* 2017;17:1–12.
- Cohen E, Davidovich M, Rokney A, Valinsky L, Rahav G, Gal-Mor O. Emergence of new variants of antibiotic resistance genomic islands among multidrug-resistant *Salmonella enterica* in poultry. *Environ Microbiol.* 2020;22:413–32.
- Aviv G, Rahav G, Gal-Mor O. Horizontal transfer of the *Salmonella enterica* serovar infantis resistance and virulence plasmid pESI to the gut microbiota of warm-blooded hosts. *MBio.* 2016;7. DOI: 10.1128/mBio.01395-16.
- Wallmann J. Antimicrobial resistance: challenges ahead. *VetRecord.* 2014;175:323–4.
- Bangera SR, Umakanth S, Chowdhury G, Saha RN, Mukhopadhyay AK, Ballal M. Poultry: a receptacle for non-typhoidal salmonellae and antimicrobial resistance. *Iran J Microbiol.* 2019;11:31–8.
- MAPA (Ministerio de Agricultura Pesca y Agricultura). Real Decreto 53/2013, de 1 de febrero, por el que se establecen las normas básicas aplicables para la protección de los animales utilizados en experimentación y otros fines científicos, incluyendo la docencia. Boletín Oficial del Estado 2013. 11370–421. <https://www.boe.es/eli/es/rd/2013/02/01/53>.
- Matuschek E, Brown DFJ, Kahlmeter G. Development of the EUCAST disk diffusion antimicrobial susceptibility testing method and its implementation in routine microbiology laboratories. *Clin Microbiol Infect.* 2014;20:O255–66.
- ISO Standard. ISO - ISO 6579-1:2017 - microbiology of the food chain — horizontal method for the detection, enumeration and serotyping of *Salmonella* — Part 1: detection of *Salmonella* spp. <https://www.iso.org/standard/56712.html>. Accessed February 2020.
- MAPA (Ministerio de Agricultura Pesca y Agricultura). Real Decreto 328 /2003, de 14 de marzo, por el que se establece y regula el plan sanitario avícola. Boletín Oficial del Estad 2003, p. 1–21. <https://www.boe.es/buscar/doc.php?id=BOE-A-2003-6802>. Accessed February 2020.
- ISO Standard. ISO - ISO/TS 6579-2:2012 - microbiology of food and animal feed — horizontal method for the detection, enumeration and serotyping of *Salmonella* — Part 2: enumeration by a miniaturized most probable number technique. <https://www.iso.org/standard/56713.html>. Accessed February 2020.
- Jarvis B, Wilrich C, Wilrich PT. Reconsideration of the derivation of Most Probable Numbers, their standard deviations, confidence bounds and rarity values. *J Appl Microbiol.* 2010;109:1660–7.
- Grimont PAD, Weill F-X. Antigenic formulae of the *Salmonella* serovars. WHO Collaborating Cent Ref Res *Salmonella.* 2007;9:1–166.
- EC (European Commission). Commission Implementing Decision of 12 November 2013 on the monitoring and reporting of antimicrobial resistance in zoonotic and commensal bacteria. Brussels, Belgium: European Commission; 2013.
- EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). EU protocol for harmonised monitoring of antimicrobial resistance in

- human *Salmonella* and *Campylobacter* isolates. Stockholm: ECDC; 2016.
33. Marin C, Lainez M. *Salmonella* detection in feces during broiler rearing and after live transport to the slaughterhouse. *Poult Sci*. 2009;88:1999–2005.
 34. Cosby DE, Cox NA, Harrison MA, Wilson JL, Jeff Buhr R, Fedorka-Cray PJ. *Salmonella* and antimicrobial resistance in broilers: a review. *J Appl Poult Res*. 2015;24:408–26.
 35. EC (European Commission). Council Directive 2007/43/EC of 28 June 2007 laying down minimum rules for the protection of chickens kept for meat production (Text with EEA relevance). Brussels, Belgium: European Commission; 2007.
 36. Scanes CG. Biology of stress in poultry with emphasis on glucocorticoids and the heterophil to lymphocyte ratio. *Poult Sci*. 2016;95:2208–15.
 37. Calefi AS, Quinteiro-Filho WM, Ferreira AJP, Palermo-Neto J. Neuroimmunomodulation and heat stress in poultry. *Worlds Poult Sci J*. 2017;73:493–504.
 38. Farag MR, Alagawany M. Physiological alterations of poultry to the high environmental temperature. *J Therm Biol*. 2018;76:101–6.
 39. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control), Koutsoumanis K, Allende A, Alvarez-Ordóñez A. Opinion of the Scientific Panel on biological hazards (BIOHAZ) on the *Salmonella* control in poultry flocks and its public health impact.pdf. *EFSA J*. 2019;17:5596.
 40. Velasquez CG, MacKlin KS, Kumar S, Bailey M, Ebner PE, Oliver HF, et al. Prevalence and antimicrobial resistance patterns of *Salmonella* isolated from poultry farms in southeastern United States. *Poult Sci*. 2018;97:2144–52.
 41. Pulido-Landínez M. Food safety - *Salmonella* update in broilers. *Anim Feed Sci Technol*. 2019;250:53–8.
 42. Andoh LA, Dalsgaard A, Obiri-Danso K, Newman MJ, Barco L, Olsen JE. Prevalence and antimicrobial resistance of *Salmonella* serovars isolated from poultry in Ghana. *Epidemiol Infect*. 2016;144:3288–99.
 43. Daehre K, Projahn M, Semmler T, Roesler U, Friese A. Extended-spectrum beta-lactamase-/AmpC Beta-lactamase-producing enterobacteriaceae in broiler farms: transmission dynamics at farm level. *Microb Drug Resist*. 2017;24:511–8.
 44. Projahn M, Daehre K, Roesler U, Friese A. Extended-spectrum-beta-lactamase and plasmid-encoded cephamycinase-producing enterobacteria in the broiler hatchery as a potential mode of pseudovertical transmission. *Appl Environ Microbiol*. 2017;83:e02364–16.
 45. Projahn M, Daehre K, Semmler T, Guenther S, Roesler U, Friese A. Environmental adaptation and vertical dissemination of ESBL-/pAmpC-producing *Escherichia coli* in an integrated broiler production chain in the absence of an antibiotic treatment. *Microb Biotechnol*. 2017;11:1017–26.
 46. Agyare C, Etsiapa Boamah V, Ngofi Zumbi C, Boateng Osei F. Antibiotic use in poultry production and its effects on bacterial resistance. In: Kumar Y, editor. *Antimicrobial resistance - a global threat*. London: IntechOpen; 2019.
 47. Osman KM, Kappell AD, Elhadidy M. Poultry hatcheries as potential reservoirs for antimicrobial-resistant *Escherichia coli*: A risk to public health and food safety. *Sci Rep*. 2018;8:5859.
 48. Okorafor ON, Anyanwu MU, Nwafor EO, Anosa GN, Udeg-bunam RI. Multidrug-resistant enterobacteria colonize commercial day-old broiler chicks in Nigeria. *Vet World*. 2019;12:418–23.
 49. Tripathi V, Tripathi P. Antibiotic resistance genes: an emerging environmental pollutant. In: *Perspectives in environmental toxicology*. Kesari K, editor. Switzerland: Environmental Science and Engineering; 2017. p. 183–201.
 50. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). The European Union Summary Report on Antimicrobial Resistance in zoonotic and indicator bacteria from humans, animals and food in 2017/2018. *EFSA J*. 2020;18:e06007.
 51. Allen HK, Donato J, Wang HH, Cloud-Hansen KA, Davies J, Handelsman J. Call of the wild: antibiotic resistance genes in natural environments. *Nat Rev Microbiol*. 2010;8:251–9.
 52. Bengtsson-Palme J, Kristiansson E, Larsson DGJ. Environmental factors influencing the development and spread of antibiotic resistance. *FEMS Microbiol Rev*. 2018;42:fux053.
 53. Westphal-Settele K, Konradi S, Balzer F, Schönfeld J, Schmithausen R. The environment as a reservoir for antimicrobial resistance: A growing problem for public health? *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitschutz*. 2018;61:533–42.
 54. Kern WV. Multiresistant bacteria - antibiotic prescription and antibiotics of last resort. *Dtsch Medizinische Wochenschrift*. 2018;143:643–50.
 55. WHO (World Health Organization). *Critically important antimicrobials for human medicine*, 6th ed. Geneva: World Health Organization; 2019.
 56. ESVAC (European Medicines Agency). Sales of veterinary antimicrobial agents in 30 European countries in 2016. Trends from 2010 to 2016. Eight ESVAC report. Amsterdam, Netherlands: EMA; 2018.
 57. EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control). Technical specifications on harmonised monitoring of antimicrobial resistance in zoonotic and indicator bacteria from food-producing animals and food. *EFSA J*. 2019;17:5709.

How to cite this article: Montoro-Dasi L, Villagra A, Vega S, Marin C. Influence of farm management on the dynamics of *Salmonella enterica* serovar Infantis shedding and antibiotic resistance during the growing period of broiler chickens. *Vet Rec*. 2021;e302.
<https://doi.org/10.1002/vetr.302>